

## Effective July 1, 2015 Immunization Rule Change

### Kindergarten

**Polio vaccine-** the booster (4<sup>th</sup>) dose is required on or after the 4<sup>th</sup> birthday and before entering school for the first time.

**Varicella Vaccine-** 2 doses administered at least 28 days apart.

(2<sup>nd</sup> dose before entering school for the first time)

### Seventh Grade

**Tetanus, diphtheria, pertussis (whooping cough) - Tdap-** booster must be given before entering 7<sup>th</sup> grade or 12 years of age whichever comes first.

**Meningococcal Conjugate Vaccine (MCV) – 2 doses**

One dose entering 7<sup>th</sup> grade or by the age of 12 whichever comes first

Booster dose for those entering 12<sup>th</sup> grade or 17 years of age beginning August 1, 2020

#### Immunization Schedule

Birth	Hepatitis B (HBV)
2 Months	HBV, Tdap, IPV, Hib, PCV, Rotavirus*
4 Months	Tdap, IPV, Hib, PCV, Rotavirus*
6Months	HBV, Tdap, IPV, Hib, PCV, Rotavirus*
Over 6 months	Flu*
12-18 months	Tdap, Hib, MMR, Varicella, PCV, HAV (Hepatitis A)*
4-6 years	Tdap, IPV, MMR, Varicella
7 <sup>th</sup> Grade	Tdap & Meningococcal (1 dose entering 7 <sup>th</sup> grade)

\*are recommended only Rotavirus and Hepatitis A